

1. Goetz, S. J., Sun, M., Zolkos, S., Hansen, A., & Dubayah, R. (2014). The relative importance of climate and vegetation properties on patterns of North American breeding bird species richness. *Environmental Research Letters*, 9(3), 034013. doi:10.1088/1748-9326/9/3/034013. <http://iopscience.iop.org/1748-9326/9/3/034013/article/>
2. Goetz, S.J., et al. (2010). Lidar remote sensing variables predict breeding habitat of a Neotropical migrant bird. *Ecology*, 91, 1569-1576. <http://onlinelibrary.wiley.com/doi/10.1890/09-1670.1/full>
3. Swatantran, A., Dubayah, R., Goetz, S., Hofton, M., Betts, M.G., Sun, M., Simard, M., & Holmes, R. (2012). *Mapping migratory bird prevalence using remote sensing data fusion*. PLoS ONE 7(1):e28922. <http://journals.plos.org/plosone/article?id=10.1371/journal.pone.0028922>
4. Goetz, S.J., et al. (2007). Laser remote sensing of canopy habitat heterogeneity as a predictor of bird species richness in an eastern temperate forest, USA. *Remote Sensing of Environment*, 108, 254-263. http://cce.nasa.gov/veg3dbiomass/goetz_etal2007.pdf
5. Davies, A. B., & Asner, G. P. (2014). Advances in animal ecology from 3D-LiDAR ecosystem mapping. *Trends in Ecology & Evolution*, 29(12), 681–91. doi:10.1016/j.tree.2014.10.005. www.cell.com/article/S0169534714002249/fulltext